

Ratings of Withholding Aerosol Therapies for Hemoptysis* and Pneumothorax**

Therapy	Mild-to-Moderate Hemoptysis			Massive Hemoptysis			Small Pneumothorax			Large Pneumothorax		
	Median	IQR	Consensus	Median	IQR	Consensus	Median	IQR	Consensus	Median	IQR	Consensus
All	2	2–3	Good	5	3–8	None	0	0–1	Very Good	1	0–2.25	Good
Hypertonic Saline	5	3–8	Some	8	5–9.5	Some	3	1–6	Some	4	1–7	None
Dornase alfa	4	2–6	Some	6	2–6	Some	1	0–2	Very Good	1.5	0.75–4.25	Some
Broncho-dilators	2	1–3	Good	4	1–3	Some	0.5	0–1	Very Good	1	0–2.25	Good
Antibiotics	2	1–3.5	Good	4	2–7	Some	1	0–2	Very Good	1	0–2.25	Good

*In response to the statements, “The patient with (*mild-to-moderate* or *massive*) hemoptysis should always stop (*all aerosol medications* or *specific aerosol medication*).”

**In response to the statements: “The patient with (*small* or *large*) pneumothorax should always stop (*all aerosol medications* or *specific aerosol medication*).”

IQR: interquartile range